

Dietary patterns, obesity markers and leukocyte telomere length among Brazilian civil servants: Cross-sectional results from the Pro-Saude study

Nathalia Ferrazzo Napolini^{1*}, Rosely Sichieri¹, Diana Cunha Barbosa¹, Rosangela A. Pereira², Eduardo Faerstein¹

¹Instituto de Medicina Social, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil.

²Universidade Federal do Rio de Janeiro. Departamento de Nutrição Social e Aplicada. Rio de Janeiro, RJ, Brasil

***Corresponding author:** Nathalia Ferrazzo Napolini, Institution: Instituto de Medicina Social, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil. Address: Bloco D e E, R. São Francisco Xavier, 524 - 7º andar - Maracanã, Rio de Janeiro - RJ, 20550-900, Telephone: +5521983302333, Email address: nfnapolini@usp.br

Job Title: Postdoctoral Associate

Keywords: Dietary patterns, Partial least squares, Obesity, Telomere length

Running title: Diet, obesity, and telomere length



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980023001064

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Acknowledgements

The authors are grateful to the public civil servant who participated in this study and, to the Universidade do Estado do Rio de Janeiro for their technical and logistical assistance and research support.

Financial Support

This study was supported by Grant E-26/110.119/2014 from FAPERJ (Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro)
NFN received a posdoctoral scholarship from FAPERJ (Pós-Doutorado Nota 10; 204.218/2021)

Conflict of Interest

The authors declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.

Authorship

Conceptualization: RS, RAP, DCB

Data analysis and interpretation: NFN, RS, DCB, RAP

Literature search: NFN, RS, EF

Project coordination and funding acquisition: EF, RS

Supervision: RS, RAP, DCB

Writing—original draft: NFN

Writing—review & editing: NFN, RS, DCB, RAP, EF

Ethical Standards Disclosure

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Research Ethics Committee of the Institute of Social Medicine, State of University of Rio de Janeiro (CAAE 0041.0.259.000-11). Written informed consent was obtained from all participants before data collection.

ABSTRACT

Objective: Dietary patterns express the combination and variety of foods in the diet. The Partial Least Squares (PLS) method allows extracting dietary patterns related to a specific health outcome. Few studies have evaluated obesity-related dietary patterns associated with telomeres length. This study aims to identify dietary patterns explaining obesity markers and to assess their association with leucocyte telomere length (LTL), a biological marker of the aging process.

Design: Cross-sectional study.

Setting: University campuses in the state of Rio de Janeiro, Brazil

Participants: 478 participants of a civil servants' cohort study with data on food consumption, obesity measurements (total body fat, visceral fat, BMI, leptin, adiponectin), and blood samples.

Results: Three dietary patterns were extracted: (1) fast food and meat; (2) healthy; and (3) traditional pattern, which included rice and beans, the staple foods most consumed in Brazil. All three dietary patterns explained 23.2% of food consumption variation and 10.7% of the obesity-related variables. The fast food and meat pattern was the first factor extracted, explaining 11-13% variation of the obesity-related response variables (BMI, total body fat and visceral fat), leptin and adiponectin showed the lowest percentage (4.5-0.1%). The healthy pattern mostly explained leptin and adiponectin variations (10.7 and 3.3%, respectively). The traditional pattern was associated with LTL ($\beta = 0.0117$; 95% CI 0.0001 - 0.0233) after adjustment for the other patterns, age, sex, exercise practice, income, and energy intake.

Conclusion: Leucocyte telomere length was longer among participants eating a traditional dietary pattern that combines fruit, vegetables, and beans.

Abbreviations: BMI: Body mass index, COVID-19: Coronavirus disease, FFQ: Food Frequency Questionnaire, LTL: Leukocyte telomere length, NADPH: Nicotinamide adenine dinucleotide phosphate, NCDs: Non-communicable chronic diseases, PLS: Partial Least Squares

INTRODUCTION

Over the past four decades, the global prevalence of obesity has nearly tripled ⁽¹⁾, making it a global pandemic ⁽²⁾. In Brazil, the prevalence of obesity in adults increased by 54% from 2003 to 2019 and the prevalence of overweight increased by 30% in the same period ⁽³⁾. Obesity increases the risk of many non-communicable chronic diseases (NCDs) ⁽⁴⁾. In 2016, an estimated 5 million deaths were attributed to overweight and obesity worldwide ⁽⁵⁾.

The mechanisms linking obesity to many NCDs encompass dysfunctional adipose tissue inducing systemic low-grade inflammation and oxidative stress ^(6, 7). The increased levels of proinflammatory cytokines and acute-phase proteins are related to the secretory capacity of the adipose tissue and its dysfunctional characteristic in this condition. Systemic inflammation linked to obesity seems to be determinant for the development of several comorbidities, such as insulin resistance, type 2 diabetes mellitus, and cardiovascular diseases ⁽⁸⁾. Telomere length may be influenced by obesity-related inflammation and oxidative stress ⁽⁹⁾.

Telomeres are regions of repeated nucleotide sequences (TTAGGG in humans), situated at both ends of each chromosome and protect against chromosome degradation and inter-chromosomal fusion. Telomeres become gradually shorter with increasing age due to DNA loss during cell division, representing a biologic marker of aging ⁽¹⁰⁾. A meta-analysis of 87 observational studies found an inverse association between body mass index and telomere length ⁽¹¹⁾. Yet, no prior studies have focused on the impact of diet and dietary patterns associated with obesity that may modify the telomeres. The most consistent association found by reviews was with Mediterranean diet, with higher adherence to this dietary pattern associated with longer telomeres ^(12, 13).

Exploratory dietary pattern analyses could realistically reflect dietary habits by examining the effects of the total diet. The PLS method combines an exploratory analysis, with a priori hypothesis on the role of intermediate variables on the outcome. This hybrid methodologies have gained interest as unique data reduction techniques for establishing a direct link between dietary exposures and health outcomes ⁽¹⁴⁾.

Nevertheless, studies evaluating dietary patterns explaining obesity-related markers and their association with telomeres length have been poorly described. In the present study, we used the PLS method to identify dietary patterns explaining the following obesity-related markers: body fat, visceral fat, BMI, leptin, adiponectin, and to further assess their association with LTL.

METHODS

Study design and population

A cross-sectional analysis was conducted on the Pró-Saúde Study, a civil servant-based cohort study at university campuses in the state of Rio de Janeiro, Brazil. Briefly, four waves of data collection were conducted among 3253 participants (1999, 2001, 2006, 2012). During wave four (2012), a random sample of 520 participants stratified by sex, age (<50 vs. >50 years) and education level (<high school vs. >high school) from the Pró-Saúde baseline population was selected to perform additional interview, anthropometric and body composition assessment, and collection of blood samples. Details on the study design and cohort profile can be found elsewhere ⁽¹⁵⁾. Individuals with missing data for obesity measurements and diet (N=42) were excluded. Therefore, the current analysis includes data from 478 participants.

All field workers were trained, and all measurements underwent strict quality control with respect to the data collection process. In parallel with data collection, questionnaires were evaluated and coded by field supervisors, independently entered by 2 research assistants, and stored in EpiData 3.1 (EpiData Association, Odense, Denmark). Reporting of clinical data complies to the STROBE guidelines.

Diet assessment

Food consumption was assessed by using a self-reported semi-quantitative Food Frequency Questionnaire, validated by Sichieri et al. ⁽¹⁶⁾, structured on 82 food items or food groups. Food amounts are predefined and expressed as household measures or food units. Respondents selected from eight consumption frequencies (> 3 times/day; 2–3 times/day; 1 time/day; 5–6 times/week; 2–4 times/week; 1 time/week; 1–3 times month; never or almost never), converted into grams per day. The nutrient and calorie contents of the foods was derived from the Brazilian Table of Food Compositions ⁽¹⁷⁾. The usual portion/serving size

was used to estimate the daily consumption. The 82-item FFQ were assigned into 23 food groups (Suppl. Table 1) according to composition. The total dietary energy was calculated and to reduce spurious variability the dietary energy density was estimated by dividing food amounts (grams) by the daily energy consumed (kilocalories) ⁽¹⁸⁾.

Adiposity assessment

The total body mass, total body fat, and visceral fat were measured via DXA with iDXA Lunar equipment (GE Healthcare, WI) and enCORE 2008 version 12.20 software ⁽¹⁹⁾. For the full-body examination, participants wore light clothing and put off metal accessories. Participants were placed in the dorsal recumbent position and asked to remain motionless until the end of the procedure. All scans were performed by the same trained professional, and the equipment was calibrated daily according to the manufacturer's protocol. Measurements of the manufacturer-supplied calibration block (daily) and column calibration block (weekly) showed a <0.7 % coefficient of variation ⁽¹⁹⁾. The total body fat was expressed in kg and as a percentage. Visceral fat was expressed as a percentage of the total fat mass obtained via DXA.

Body mass (kg) and height (m) were measured by trained research assistants, using a digital scale with accuracy of 0.1 kg and a fixed stadiometer (SECA) with accuracy of 0.1 cm. Body mass index ($BMI = \text{body mass}/\text{height}^2$) of each participant was calculated (kg/m^2).

Blood collection and biochemical biomarkers assessment

Blood samples were collected after 12-h overnight fast by a trained professional, using vacutainer tubes (Becton, Dickinson & Company, Brazil). Serum samples were collected and stored at -80 °C until analyses. Plasma leptin (Cat. #ZHL-80SK) and adiponectin (Cat. #EZHADP-61K) levels were determined using semiautomated chemiluminescent enzyme-labeled immunometric assay (Liaison, Diasorin, MN, USA). Intra-assay coefficients of variation were < 6%.

Leukocyte telomere length assessment (LTL)

DNA samples were isolated from whole blood using a commercial kit (Puregene Blood Kit Qiagen, Hilden, Germany). The quality and concentration of DNA samples were evaluated by spectrophotometry (BioDrop DUO, BioDrop, Cambridge, England) and were stored at -80°C, until the determination of LTL and genotyping. LTL was measured by quantitative real-time polymerase chain reaction (qPCR) based on the method described by

Cawthon with modifications^(10, 20, 21). Reactions were conducted in triplicate and included genomic DNA (1.6 ng), 2 x RotorGene SYBR Green, PCR Master Mix (Qiagen, Hilden, Germany), primers Tel Forward (300 nM -CGGTTTGTGGTTGGGTTGGGTTGGGTTGGGTT) and Tel Reverse (300 nM -GGCTTGCCTTACCCTTACCCTTACCCTTACCCTTACCCT) or primer single gene Hbg1 Forward (300 nM -GCTTCTGACACA ACTGTGTTCACTAGC) and single gene Hbg2 Reverse (700 nM -CACCAACTTCATCCACGTTCAACC) (Integrated DNATechnologies, Coralville, IA, USA), in a 24 mL reaction. Amplification was conducted in a Rotor-Gene Q Real-Time PCR cycler (Qiagen, Hilden, Germany) as follows: 95°C by 5 min, 25 cycles (telomere reaction) or 35 cycles (single gene reaction) at 98°C by 7s, and 60°C by 10s. The telomere length for each sample was determined using the telomere-to-single copy gene ratio (T/S ratio) by the ΔC_t method ($C_t[\text{telomere}]/C_t[\text{single gene}]$). The T/S ratio for each sample (x) was normalized to the mean a T/S ratio of reference sample (r) [$2^{(\Delta C_{tx} - \Delta C_{tr})} = 2^{-\Delta \Delta C_t}$], which was also used for the standard curve, both as a reference sample and as an interassay validation sample. All sample data included had reference samples with an interassay coefficient of variation (CV) <9%. For qPCR validation purposes, the LTL was evaluated in a random sample (10%) using the Southern Blot method of Terminal restriction fragment (TRF), as described by Gutierrez-Rodriguez et al.⁽²²⁾. The analysis was conducted using a commercial kit (Telo TAGGG Telomere Length Assay Roche Applied Science, Indianapolis, IN, USA).

Covariates

Information on sociodemographic variables was collected using a self-report questionnaire. Variables considered in this study were: sex (male/female), age, *per capita* family income, and energy intake (kcal/day), and exercise practice. The last was assessed by means of a yes/no question: “In the last 2 weeks, did you engage in any physical activity to improve your health and physical condition or for fitness or leisure purposes?”.

Statistical analyses

Characteristics of participants were described as means and standard deviations, for continuous variables, or as proportions (%) for categorical variables. PLS method was used to derive dietary patterns that explained maximum variation in obesity-related response variables: total body fat (kg), visceral fat (kg), BMI (kg/m^2), leptin (ng/mL) and, adiponectin

(ng/mL). The response variables used in this study were chosen a priori and used as proxy for obesity and mediators of the association between dietary patterns and LTL^(11,23).

Based on the cumulative percentage of explained variance and interpretability of the dietary patterns, three factors were set to be retained in the models. Food groups having factor loadings $> |0.15|$ were considered relevant in the definition of the dietary patterns. Subsequent regression analyses were performed to estimate the association between the dietary patterns factors scores and LTL. Potential confounding effects of age, sex, exercise practice, income, and energy intake were adjusted in the model. The significance level for all analysis was set at 5%. All analyses were conducted in SAS (SAS Institute Inc., Cary, NC, USA).

Ethics approval and consent to participate

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Research Ethics Committee of the Institute of Social Medicine, State of University of Rio de Janeiro (CAAE 0041.0.259.000-11). Written informed consent was obtained from all participants before data collection.

RESULTS

The sub-sample of 478 participants from the Pró-Saúde Study, does not differ at the baseline from the overall cohort in relation to age, exercise practice, and obesity. Individuals aged 33-79 years (mean: 51.6 ± 7.9 years), with 52.3% women. Prevalence of obesity was 32.5% and about 40% were physically active (Table 1).

Factor loadings of the first three dietary patterns derived by PLS method are shown in Table 2. Food groups with absolute values of loadings ≥ 0.15 are highlighted. The first pattern was called “fast food and meat”, consisting of soft drinks, cold cuts, fast foods and savory snacks, alcoholic beverages, and red meat as positive contributors while sugar, beans, roots, pulses, milk and dairy, cake and pastries were inverse contributors. The second pattern, called “healthy” pattern had positive loadings for milk and dairy, fresh fruits, vegetables and greens, caffeinated drinks, pulses, fruit juices, and fish and negative loadings on beans, rice, pasta, alcoholic beverages, red meat, and soft drinks. The “traditional” pattern had positive loadings for usually consumed staple foods such as rice, beans, vegetables, and greens. Fruits,

fish, and fruit juices were also included in this pattern, while negative loadings were estimated for cake and pastries and fast food and savory snacks (Table 2).

The three dietary patterns together explained 23.2% of the variation of the predictive variables (food groups/items) and 10.7% of the response variables variance. Among the response variables, 14.1% of total body fat is explained by the variation in food consumption. The other anthropometric measures had similar percentage values (12.3% for BMI and 11.6% for visceral fat), whereas adiponectin showed the lowest percentage (4.7%).

The fast food and meat dietary pattern explained 13% of total body fat, 7.1% of visceral fat, and 11.4% of BMI variations, explaining 7.2% of the response variables in total. It also explained 4.5% of leptin and 0.1% of adiponectin variations. The healthy pattern mostly explained leptin and adiponectin variations (10.7 and 3.3%, respectively) (Table 2).

The traditional dietary pattern was directly associated with LTL ($\beta = 0.0117$; 95% CI 0.0001 - 0.0233) in a model including all three patterns and adjusted for age, sex, physical activity, income, and energy intake (Table 3). Separate models for each pattern showed similar results (Table 3).

DISCUSSION

The present study identified three dietary patterns related to obesity markers: “fast food and meat”, “healthy”, and “traditional”. The fast food and meat pattern explained most of the variation of the obesity-related response variables and the traditional pattern was directly associated with LTL independent of sex, age, exercise practice, income, and energy intake.

Fast foods are energy-dense, high-fat, and low-fiber content and the consumption of these foods has been associated with obesity worldwide⁽²⁴⁻²⁶⁾. This pattern of diet is usually identified as western and in a study in Iran, it was directly associated with BMI and fat mass, whereas traditional pattern was inversely associated with dietary inflammatory index and insulin resistance among healthy adults with obesity⁽²⁷⁾. This corroborates our findings, showing that the major dietary pattern responsible for explaining the obesity-related variables was the fast food and meat pattern, responsible for 7% of the response variables variance.

In contrast, healthy patterns have been inversely associated with BMI, waist circumference and other obesity-related biomarkers^(25, 28). In Brazil, the traditional dietary pattern, which is considered a healthy one and is heavily based on the consumption of rice

and beans. This pattern was protective for obesity in studies with adults^(29, 30) and adolescents^(31, 32). Authors have attributed this protective effect to the high fiber content of rice and beans combination resulting in low glycemic index meals⁽³⁰⁾. Besides, the consumption of these foods seems to be a marker of home-cooked meals that are more likely to contain vegetables and whole grains⁽³³⁾. Also, the consumption of black bean, the most consumed bean in Rio de Janeiro, is associated with reduced percentage of body fat, serum leptin, glucose and insulin concentrations, results that could be mediated in part by modification of the gut microbiota⁽³⁴⁾.

Regarding the adipokines evaluated in this study, leptin and adiponectin have important role in regulating inflammation, especially in obesity⁽⁸⁾. We found that the healthy and traditional patterns were responsible for explaining the variation of leptin (10.7% for both). Leptin is primarily known as a satiety hormone and a pro-inflammatory adipokine, its serum levels and gene expression in adipocytes strongly correlate with the proportion of body fat stores⁽⁸⁾. The close values for the explanation of leptin by the healthy and traditional patterns may be due to the high prevalence of overweight and obesity in this population. On the other hand, adiponectin's variation was mostly explained (4.7%) by the traditional pattern. Adiponectin is an anti-inflammatory factor and its production is frequently reduced in obesity⁽³⁵⁾. However, in other studies, adiponectin's levels are mostly correlated to "Western-type" and meat-based patterns and ultra-processed food consumption^(36, 37), and the later association was mediated by the presence of obesity. This divergence could be related to the dual activity of adiponectin, which acts as an anti-inflammatory factor especially in atherosclerosis, but in some chronic inflammatory/autoimmune diseases adiponectin may have pro-inflammatory effects⁽⁸⁾. Even being responsible for explaining, in part, the variation of these adipokines, the traditional patterns did not add up to the explanation of the obesity-related variables.

As observed in many countries, including Brazil⁽³⁸⁾ there is a trend of reduction in the consumption of traditional foods, and substitution by ultra-processed foods. High intake of ultra-processed foods has been associated with a higher risk of having shorter telomeres in an elderly Spanish population⁽³⁹⁾. In contrast, the traditional dietary pattern in Costa Rica – which also has rice and beans as staple food – was directly associated with telomere length in an elderly population⁽⁴⁰⁾. A systematic review and meta-analysis of eight cross-sectional studies demonstrated that higher Mediterranean diet adherence is associated with longer telomere⁽¹²⁾. The Mediterranean diet is a traditional dietary patter in countries such as

Greece, Spain, and Italy in the period of pre-globalization of the food system. Another narrative review of Mediterranean diet demonstrated its positive influence on longevity⁽¹³⁾. Moreover, telomere length was linearly related to fruits and vegetables intake in a study of 5,448 U.S. adults⁽⁴¹⁾.

The traditional dietary pattern identified in our analysis has common characteristics with the Mediterranean dietary pattern, such as high consumption of vegetables including green vegetables, fruits, and beans (legumes/pulses) and, moderate intakes of fish, and low intakes of sweets; and differs regarding the consumption of extra virgin olive oil, nuts, dairy products, and red wine⁽⁴²⁾ – which are not usually consumed. Beyond other positive aspects of a traditional food intake, such as reduction in weight gain, our results indicate a role in longevity reinforcing dietary guidelines of increasing intake of the traditional Brazilian dietary pattern. Together with the positive findings in other countries, our results consolidate a protective role for traditional dietary patterns.

The limitation of the study is the cross-sectional analysis of the cohort since the LTL measure and food consumption was done after 13 years of follow-up. This may explain the similar estimate values, although in opposite direction, for healthy and traditional patterns. It might represent a higher adherence to the healthy pattern among individuals with higher BMI since reverse causality cannot be ruled out. Furthermore, our study sample of civil servants limits the possibility to extrapolate results to the general population. However, we showed that the subsample with LTL measures had similar distribution by age, BMI and exercise practice compared to the baseline.

This study also has strengths, particularly the use of the PLS to identify eating patterns. The PLS is a hybrid method that extracts a latent factor simultaneously related to a set of independent variables and to a set of dependent variables, in such way that the explanation of the covariance between the two set of variables is maximized and a direct link between dietary exposures and clinical outcomes is established. In this case, data reduction of food groups is directed by biomarkers of interest. Therefore, this technique may be more suitable to extract dietary patterns when the objective is to explain the effect of intermediary variables in the relationship of eating patterns and an outcome.

Our results indicate that the consumption of traditional and culturally acceptable foods must be encouraged to promote health, longevity, and control obesity.

References

1. NCD-RisC. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies with 128.9 million participants.: *Lancet*; 2017.
2. Swinburn BA, Kraak VI, Allender S, Atkins VJ, Baker PI, Bogard JR, et al. The Global Syndemic of Obesity, Undernutrition, and Climate Change: The Lancet Commission report. *Lancet*. 2019;393(10173):791-846.
3. PNS. Pesquisa Nacional de Saúde. Atenção primária à saúde e informações antropométricas: Brasil/2019. Rio de Janeiro: IBGE, Coordenação de Trabalho e Rendimento 2020. p. 66p.
4. WHO. Noncommunicable diseases country profiles 2018. Geneva: World Health Organization 2018.
5. Collaborators GRF. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016.: *Lancet*; 2017. p. 1345–422.
6. Fernández-Sánchez A, Madrigal-Santillán E, Bautista M, Esquivel-Soto J, Morales-González A, Esquivel-Chirino C, et al. Inflammation, oxidative stress, and obesity. *Int J Mol Sci*. 2011;12(5):3117-32.
7. Wu H, Ballantyne CM. Metabolic Inflammation and Insulin Resistance in Obesity. *Circ Res*. 2020;126(11):1549-64.
8. Aprahamian TR, Sam F. Adiponectin in cardiovascular inflammation and obesity. *Int J Inflam*. 2011;2011:376909.
9. Zhou Y, Hambly BD, McLachlan CS. FTO associations with obesity and telomere length. *J Biomed Sci*. 2017;24(1):65.
10. Lin J, Epel E, Cheon J, Kroenke C, Sinclair E, Bigos M, et al. Analyses and comparisons of telomerase activity and telomere length in human T and B cells: insights for epidemiology of telomere maintenance. *J Immunol Methods*. 2010;352(1-2):71-80.

11. Gielen M, Hageman GJ, Antoniou EE, Nordfjall K, Mangino M, Balasubramanyam M, et al. Body mass index is negatively associated with telomere length: a collaborative cross-sectional meta-analysis of 87 observational studies. *Am J Clin Nutr.* 2018;108(3):453-75.
12. Canudas S, Becerra-Tomás N, Hernández-Alonso P, Galié S, Leung C, Crous-Bou M, et al. Mediterranean Diet and Telomere Length: A Systematic Review and Meta-Analysis. *Adv Nutr.* 2020;11(6):1544-54.
13. Shannon OM, Ashor AW, Scialo F, Saretzki G, Martin-Ruiz C, Lara J, et al. Mediterranean diet and the hallmarks of ageing. *Eur J Clin Nutr.* 2021;75(8):1176-92.
14. Zhao J, Li Z, Gao Q, Zhao H, Chen S, Huang L, et al. A review of statistical methods for dietary pattern analysis. *Nutr J.* 2021;20(1):37.
15. Nery Guimarães JM, Chor D, Werneck GL, Carvalho MS, Coeli CM, Lopes CS, et al. Association between self-rated health and mortality: 10 years follow-up to the Pró-Saúde cohort study. *BMC Public Health.* 2012;12:676.
16. Sichieri R, Everhart E. Validity of a Brazilian food frequency questionnaire against dietary recalls and estimated energy intake. *Nutrition Research;* 1998. p. 1649-59.
17. Brasil. Tabela Brasileira de Composição de Alimentos (TACO). 1ª ed. ed. Campinas: NEPA – UNICAMP; 2004.
18. Willett WC, Stampfer M, Tobias DK. Re: Adjustment for energy intake in nutritional research: a causal inference perspective. *Am J Clin Nutr.* 2022.
19. GE i. Medical Systems enCORE Operator's Manual and Product Information. Madison, WI: Lunar Corp.
20. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res.* 2002;30(10):e47.
21. Calado RT, Cooper JN, Padilla-Nash HM, Sloand EM, Wu CO, Scheinberg P, et al. Short telomeres result in chromosomal instability in hematopoietic cells and precede malignant evolution in human aplastic anemia. *Leukemia.* 2012;26(4):700-7.

22. Gutierrez-Rodrigues F, Santana-Lemos BA, Scheucher PS, Alves-Paiva RM, Calado RT. Direct comparison of flow-FISH and qPCR as diagnostic tests for telomere length measurement in humans. *PLoS One*. 2014;9(11):e113747.
23. Jessri M, Wolfinger RD, Lou WY, L'Abbé MR. Identification of dietary patterns associated with obesity in a nationally representative survey of Canadian adults: application of a priori, hybrid, and simplified dietary pattern techniques. *Am J Clin Nutr*. 2017;105(3):669-84.
24. Elizabeth L, Machado P, Zinöcker M, Baker P, Lawrence M. Ultra-Processed Foods and Health Outcomes: A Narrative Review. *Nutrients*. 2020;12(7).
25. Schlesinger S, Neuenschwander M, Schwedhelm C, Hoffmann G, Bechthold A, Boeing H, et al. Food Groups and Risk of Overweight, Obesity, and Weight Gain: A Systematic Review and Dose-Response Meta-Analysis of Prospective Studies. *Adv Nutr*. 2019;10(2):205-18.
26. Dicken SJ, Batterham RL. The Role of Diet Quality in Mediating the Association between Ultra-Processed Food Intake, Obesity and Health-Related Outcomes: A Review of Prospective Cohort Studies. *Nutrients*. 2021;14(1).
27. Saghafi-Asl M, Mirmajidi S, Asghari Jafarabadi M, Vahid F, Shivappa N, Hébert JR, et al. The association of dietary patterns with dietary inflammatory index, systemic inflammation, and insulin resistance, in apparently healthy individuals with obesity. *Sci Rep*. 2021;11(1):7515.
28. Beck KL, Jones B, Ullah I, McNaughton SA, Haslett SJ, Stonehouse W. Associations between dietary patterns, socio-demographic factors and anthropometric measurements in adult New Zealanders: an analysis of data from the 2008/09 New Zealand Adult Nutrition Survey. *Eur J Nutr*. 2018;57(4):1421-33.
29. Sichieri R, Castro JF, Moura A. [Factors associated with dietary patterns in the urban Brazilian population]. *Cad Saude Publica*. 2003;19 Suppl 1:S47-53.
30. Cunha DB, de Almeida RM, Sichieri R, Pereira RA. Association of dietary patterns with BMI and waist circumference in a low-income neighbourhood in Brazil. *Br J Nutr*. 2010;104(6):908-13.

31. Rocha NP, Milagres LC, Filgueiras MS, Suhett LG, Silva MA, Albuquerque FM, et al. Association of Dietary Patterns with Excess Weight and Body Adiposity in Brazilian Children: The Pase-Brasil Study. *Arq Bras Cardiol.* 2019;113(1):52-9.
32. Rodrigues PR, Pereira RA, Cunha DB, Sichieri R, Ferreira MG, Vilela AA, et al. Factors associated with dietary patterns in adolescents: a school-based study in Cuiabá, Mato Grosso. *Rev Bras Epidemiol.* 2012;15(3):662-74.
33. Fertig AR, Loth KA, Trofholz AC, Tate AD, Miner M, Neumark-Sztainer D, et al. Compared to Pre-prepared Meals, Fully and Partly Home-Cooked Meals in Diverse Families with Young Children Are More Likely to Include Nutritious Ingredients. *J Acad Nutr Diet.* 2019;119(5):818-30.
34. Sánchez-Tapia M, Hernández-Velázquez I, Pichardo-Ontiveros E, Granados-Portillo O, Gálvez A, R Tovar A, et al. Consumption of Cooked Black Beans Stimulates a Cluster of Some Clostridia Class Bacteria Decreasing Inflammatory Response and Improving Insulin Sensitivity. *Nutrients.* 2020;12(4).
35. Stofkova A. Leptin and adiponectin: from energy and metabolic dysbalance to inflammation and autoimmunity. *Endocr Regul.* 2009;43(4):157-68.
36. Barbaresko J, Koch M, Schulze MB, Nöthlings U. Dietary pattern analysis and biomarkers of low-grade inflammation: a systematic literature review. *Nutr Rev.* 2013;71(8):511-27.
37. Lopes AEDS, Araújo LF, Levy RB, Barreto SM, Giatti L. Association between consumption of ultra-processed foods and serum C-reactive protein levels: cross-sectional results from the ELSA-Brasil study. *Sao Paulo Med J.* 2019;137(2):169-76.
38. Rodrigues RM, Souza AM, Bezerra IN, Pereira RA, Yokoo EM, Sichieri R. Most consumed foods in Brazil: evolution between 2008-2009 and 2017-2018. *Rev Saude Publica.* 2021;55(Supl 1):4s.
39. Alonso-Pedrero L, Ojeda-Rodríguez A, Martínez-González MA, Zalba G, Bes-Rastrollo M, Marti A. Ultra-processed food consumption and the risk of short telomeres in an elderly population of the Seguimiento Universidad de Navarra (SUN) Project. *Am J Clin Nutr.* 2020;111(6):1259-66.

40. Ruiz-Narváez EA, Baylin A, Azofeifa J, Leal A, Rosero-Bixby L. Diet and Leukocyte Telomere Length in a Population with Extended Longevity: The Costa Rican Longevity and Healthy Aging Study (CRELES). *Nutrients*. 2021;13(8).
41. Tucker LA. Fruit and Vegetable Intake and Telomere Length in a Random Sample of 5448 U.S. Adults. *Nutrients*. 2021;13(5).
42. Davis C, Bryan J, Hodgson J, Murphy K. Definition of the Mediterranean Diet; a Literature Review. *Nutrients*. 2015;7(11):9139-53.

Table 1 – Sample size, means, percentages and standard deviation (SD) of the study population. Pro-Saúde Study—Rio de Janeiro, Brazil, 2012–2013.

	N	mean	SD
Age, years	478	51.7	7.7
Body mass index (kg/m ²)	478	28.1	5.1
Height (cm)	478	165.9	9.2
Total body lean mass (kg)	478	46.1	9.8
Total body fat (kg)	478	28.1	9.9
Visceral fat (kg)	478	13.2	8.9
Energy intake, kcal/day	478	2487.2	922.1
Leucocyte telomere length, db	478	0.58	0.15
Leptin	478	6.3	5.7
Adiponectin	478	53.4	46.5
	N	%	
Sex			
Male	228	47.7	
Female	250	52.3	
BMI categories^a			
Underweight	4	0.8	
Healthy weight	132	27.6	
Overweight	188	39.3	
Obesity	154	32.3	
Physically active^b	194	40.6	

^aBody mass index < 18.5 kg/m²: underweight; 18.5 - 24.9 kg/m²: healthy weight; 25 - 29.9 kg/m²: overweight; > 30 kg/m²: obesity

^bAny exercise practice in the last two weeks

Table 2 – Food items and groups, model effect loadings and explained variance (%) from dietary patterns associated to obesity derived from partial least squares (PLS).

Food items/groups	Model Effect Loadings		
	1 “Fast food and Meat”	2 “Healthy”	3 “Traditional”
Soft drinks	0.435	-0.177	-0.110
Cold cuts	0.406	-0.030	-0.111
Sugar	-0.340	-0.125	-0.141
Beans	-0.318	-0.348	0.277
Fast foods, savory snacks	0.303	-0.088	-0.329
Roots	-0.232	-0.077	0.009
Alcoholic beverages	0.213	-0.244	0.088
Red meat	0.209	-0.209	-0.117
Pulses	-0.199	0.179	0.092
Milk and dairy	-0.197	0.356	0.082
Cake and pastries	-0.195	0.098	-0.41
Rice	-0.105	-0.343	0.409
Fresh fruits	-0.101	0.348	0.282
Fish	0.100	0.168	0.256
Caffeinated drinks	0.064	0.213	0.117
Vegetables and greens	0.039	0.326	0.391
Pasta	-0.017	-0.280	-0.073
Fruit juices	0.016	0.169	0.224
Cumulate total explained variation (%)	6.71	15.17	23.22
Response Variables	% variation accounted for		
Leptin	4.55	10.67	10.68
Adiponectin	0.06	3.27	4.69
Total body fat (kg)	13.04	13.79	14.13
Visceral fat (kg)	7.11	11.18	11.60
BMI	11.40	11.41	12.35
Cumulate total explained variation (%)	7.23	10.07	10.69

For food items/groups values are loadings and for response variables values are % variation accounted for by PLS factors.

Remaining food groups loadings had $< |0.15|$: Fats (Butter and mayonnaise), Bread, Eggs, healthy snacks (peanuts and popcorn), meats (Pork, chicken, and tripe); Number of participants included in the model=478.

Table 3 – Regression coefficients between the identified dietary patterns and telomere length. Pro-Saúde Study—Rio de Janeiro, Brazil, 2012–2013.

	Crude β	95% CI	Adjusted ¹ β	95% CI	Adjusted ² β	95% CI
Models including all three dietary patterns						
Fast food and Meat	0.0069	-0.0043 - 0.0181	0.0016	-0.0096 - 0.0128	0.0067	-0.0153 - 0.0287
Healthy	-0.0095	-0.0196 - 0.0066	-0.0108	-0.0243 - 0.0027	-0.0086	-0.0268 - 0.0096
Traditional	0.0085	-0.0017 - 0.0187	0.0107	0.0003 - 0.0211	0.0117	0.0001 - 0.0233
Models for each pattern						
	Crude β	95% CI	Adjusted ¹ β	95% CI		
Fast food and Meat	0.0024	-0.0060 - 0.0108	-0.0027	-0.0121 - 0.0067	-0.0045	-0.0167 - 0.0077
Healthy	-0.0043	-0.0143 - 0.005	-0.0101	-0.0213 - 0.0011	-0.0046	-0.0148 - 0.0056
Traditional	0.0094	-0.0008 - 0.0196	0.0112	0.0010 - 0.0214	0.0109	0.0001 - 0.0219

¹Adjusted for age, sex and, exercise practice

¹Adjusted for age, sex, exercise practice and energy intake